


Modern Numerical Continuation Methods for Biological Systems

Christian Kuehn

Lichtenberg Professor for
Multiscale and Stochastic Dynamics
Technical University of Munich (TUM)

External Faculty, Complexity Science Hub Vienna

www.multiscale.systems



Research Area(s)

Multiscale Methods

Stochastic Systems

Nonlinear Dynamical Systems

Nonlocality & Patterns

Network Dynamics

Research Area(s)

Multiscale Methods

- ▶ fast-slow systems
- ▶ perturbation methods
- ▶ geometric desingularization
- ▶ complex oscillations
- ▶ ...

Nonlocality & Patterns

- ▶ fractional & nonlocal PDEs
- ▶ numerical continuation
- ▶ travelling waves
- ▶ bifurcation theory
- ▶ ...

Stochastic Systems

- ▶ path-based methods
- ▶ early-warning signs
- ▶ stochastic PDEs
- ▶ (rigorous) computation
- ▶ ...

Network Dynamics

- ▶ adaptive networks
- ▶ graph limits
- ▶ data analysis
- ▶ moment closure
- ▶ ...

Motivation & Problem Formulation

Consider the general differential equation

$$\frac{\partial u}{\partial t} = a(u; p),$$

$p \in \mathbb{R}^n$ are parameters.

→ $a(u; p)$ could be ODE, DDE, PDE, SDE, SPDE, etc.

Motivation & Problem Formulation

Consider the general differential equation

$$\frac{\partial u}{\partial t} = a(u; p),$$

$p \in \mathbb{R}^n$ are parameters.


→ $a(u; p)$ could be ODE, DDE, PDE, SDE, SPDE, etc.

Problem: Simulation only the first step, expensive if we have to:

1. Simulate over initial values $u_0 \in \mathbb{R}^m$.
2. Simulate over parameter space $p \in \mathbb{R}^n$.
3. Simulate over noise realizations $\omega \in \Omega$.

Do you really understand the nonlinear dynamics from averages?

Motivation & Problem Formulation



Problem: Simulation only the first step, expensive if we have to:

1. Simulate over initial values $u_0 \in \mathbb{R}^m$.
2. Simulate over parameter space $p \in \mathbb{R}^n$.
3. Simulate over noise realizations $\omega \in \Omega$.

Do you really **understand the nonlinear dynamics** from **averages**?

Typical goal: “parameter vs. quantity of interest”-diagram.


Deterministic DEs Standard Method: Continuation

Consider the ODE


$$x' = f(x; p), \quad f : \mathbb{R}^m \times \mathbb{R} \rightarrow \mathbb{R}^m.$$

Let $(x; p) =: z$. A curve $z = \gamma(s)$ of **equilibria** satisfies

$$f(\gamma(s)) = 0. \quad (\text{note: } Df(\gamma(0))\gamma'(0) = 0)$$



(a) Prediction Step



(b) Correction Step

Important: Excellent guess from (a) for **Newton's Method** in (b).

Deterministic DEs Standard Method: Continuation

Consider the ODE

$$x' = f(x; p), \quad f : \mathbb{R}^m \times \mathbb{R} \rightarrow \mathbb{R}^m.$$


Let $(x; p) =: z$. A curve $z = \gamma(s)$ of **equilibria** satisfies

$$f(\gamma(s)) = 0. \quad (\text{note: } Df(\gamma(0))\gamma'(0) = 0)$$

Remarks on Software:

- ▶ AUTO-07p: Doedel, Champneys, Sandstede, et al
- ▶ XPP-Aut: Ermentrout
- ▶ MatCont: Meijer, Govaerts, Kuznetsov, et al
- ▶ pde2path: Uecker, Rademacher, Dohnal, et al
- ▶ ...

Example 1: SIRWS Epidemics



$$\dot{S} = -\frac{\beta}{\epsilon}SI + 2\kappa W + \xi(1 - S),$$


$$\dot{I} = \frac{\beta}{\epsilon}SI - \frac{\gamma}{\epsilon}I - \xi I,$$

$$\dot{R} = \frac{\gamma}{\epsilon}I - 2\kappa R + p \frac{\beta}{\epsilon}IW - \xi R,$$

$$\dot{W} = 2\kappa R - 2\kappa W - p \frac{\beta}{\epsilon}IW - \xi W,$$

Ref: "A geometric analysis of the SIR, SIRS and SIRWS epidemiological models", H. Jardon-Kojakhmetov, CK, A. Pugliese, M. Sensi, preprint, 2020.


Example 1: SIRWS Epidemics



Ref: "A geometric analysis of the SIR, SIRS and SIRWS epidemiological models", H. Jardon-Kojakhmetov, **CK**, A. Pugliese, M. Sensi, preprint, 2020.

Example 2: The 3D FitzHugh-Nagumo Equation


$$\begin{aligned}x_1' &= x_2, \\x_2' &= \frac{1}{5}p_2x_2 - x_1(1-x_1)(x_1-0.1) - y + p_1, \\y' &= \varepsilon(x_1 - \gamma y)\end{aligned}$$



Refs: "Homoclinic orbits of the FitzHugh-Nagumo equation: the singular limit", J. Guckenheimer and K., *Discrete and Continuous Dynamical Systems S*, Vol. 2, No. 4, pp. 851-872, 2009. // "Homoclinic orbits of the FitzHugh-Nagumo equation: bifurcations in the full system", J. Guckenheimer and K., *SIAM Journal on Applied Dynamical Systems*, Vol. 9, No. 1, pp. 138-153, 2010.

Example 3: Shigesada-Kawasaki-Teramoto (SKT)


$$\begin{aligned}\partial_t u &= \Delta((p_1 + d_{11}u + d_{12}v)u) + (r_1 - a_1u - b_1v)u, \\ \partial_t v &= \Delta((p_1 + d_{22}v + p_2u)v) + (r_2 - b_2u - a_2v)v,\end{aligned}$$



Ref: "On the influence of cross-diffusion in pattern formation", M. Breden, CK and C. Soresina, arXiv:1910.03436.

Example 3: Shigesada-Kawasaki-Teramoto (SKT)

$$\begin{aligned}\partial_t u &= \Delta((p_1 + d_{11}u + d_{12}v)u) + (r_1 - a_1u - b_1v)u, \\ \partial_t v &= \Delta((p_1 + d_{22}v + p_2u)v) + (r_2 - b_2u - a_2v)v,\end{aligned}$$



Ref: "On the influence of cross-diffusion in pattern formation", M. Breden, CK and C. Soresina, arXiv:1910.03436.

Example 4: Shigesada-Kawasaki-Teramoto (SKT)

Fast reaction SKT model ($u = u_1 + u_2$):

$$\begin{aligned}\partial_t u_1 &= p_1 \Delta u_1 + (p_2 - a_1 u - b_1 v) u_1 + \frac{1}{\varepsilon} \left(u_2 \left(1 - \frac{v}{M} \right) - u_1 \frac{v}{M} \right), \\ \partial_t u_2 &= (p_1 + d_{12} M) \Delta u_2 + (p_2 - a_1 u - b_1 v) u_2 \\ &\quad - \frac{1}{\varepsilon} \left(u_2 \left(1 - \frac{v}{M} \right) - u_1 \frac{v}{M} \right), \\ \partial_t v &= d_2 \Delta v + (r_2 - b_2(u_1 + u_2) - a_2 v) v.\end{aligned}$$


Cross-diffusion ($\varepsilon \rightarrow 0$ limit) SKT model:

$$\begin{aligned}\partial_t u &= \Delta((p_1 + d_{12} v) u) + (p_2 - a_1 u - b_1 v) u, \\ \partial_t v &= d_2 \Delta v + (r_2 - b_2 u - a_2 v) v,\end{aligned}$$

Ref: “Numerical continuation for a fast-reaction system and its cross-diffusion limit”,
CK and C. Soresina, SN Partial Differential Equations and Applications, accepted / to appear, 2020.


Example 4: Shigesada-Kawasaki-Teramoto (SKT)

Fast reaction SKT model bifurcation diagram:




Example 4: Shigesada-Kawasaki-Teramoto (SKT)

Fast reaction SKT model bifurcation diagram:




Example 4: Shigesada-Kawasaki-Teramoto (SKT)

Fast reaction SKT model bifurcation diagram:




Example 4: Shigesada-Kawasaki-Teramoto (SKT)

Cross-diffusion SKT model bifurcation diagram:




Example 4: Shigesada-Kawasaki-Teramoto (SKT)

Fast reaction SKT model bifurcation diagrams...




Example 4: Shigesada-Kawasaki-Teramoto (SKT)

Fast reaction SKT model bifurcation diagrams...




Example 4: Shigesada-Kawasaki-Teramoto (SKT)

Fast reaction SKT model bifurcation diagrams...




Example 4: Shigesada-Kawasaki-Teramoto (SKT)

Fast reaction SKT model bifurcation diagrams...




Example 4: Shigesada-Kawasaki-Teramoto (SKT)

Fast reaction SKT model bifurcation diagrams...



Example 4: Shigesada-Kawasaki-Teramoto (SKT)

Fast reaction SKT model bifurcation diagrams...



Numerical Bifurcation Analysis for Stochastic Systems?

Consider the **stochastic (ordinary) differential equation (SDE)**

$$dx_t = f(x_t; p) dt + \sigma F(x_t; p) dW_t, \quad x_t \in \mathbb{R}^n,$$

$W_t = (W_{1,t}, W_{2,t}, \dots, W_{k,t})^\top$ iid **Brownian motions**,
 $F(x_t; p) \in \mathbb{R}^{n \times k}$; let $\mathfrak{D}(x; p) := \sigma^2 F(x; p) F(x; p)^\top$.

Numerical Bifurcation Analysis for Stochastic Systems?

Consider the **stochastic (ordinary) differential equation (SDE)**

$$dx_t = f(x_t; \boldsymbol{p}) dt + \sigma F(x_t; \boldsymbol{p}) dW_t, \quad x_t \in \mathbb{R}^n,$$

$W_t = (W_{1,t}, W_{2,t}, \dots, W_{k,t})^\top$ iid **Brownian motions**,
 $F(x_t; \boldsymbol{p}) \in \mathbb{R}^{n \times k}$; let $\mathfrak{D}(x; \boldsymbol{p}) := \sigma^2 F(x; \boldsymbol{p}) F(x; \boldsymbol{p})^\top$.

- ▶ **Approach 1:** Forward **Monte-Carlo** simulation.
- ▶ **Problem:** **Sampling** often prohibitive (\rightarrow **gsMC**).

Numerical Bifurcation Analysis for Stochastic Systems?

Consider the **stochastic (ordinary) differential equation (SDE)**

$$dx_t = f(x_t; \mathbf{p}) dt + \sigma F(x_t; \mathbf{p}) dW_t, \quad x_t \in \mathbb{R}^n,$$

$W_t = (W_{1,t}, W_{2,t}, \dots, W_{k,t})^\top$ iid **Brownian motions**,
 $F(x_t; \mathbf{p}) \in \mathbb{R}^{n \times k}$; let $\mathfrak{D}(x; \mathbf{p}) := \sigma^2 F(x; \mathbf{p}) F(x; \mathbf{p})^\top$.

- ▶ **Approach 1:** Forward **Monte-Carlo** simulation.
- ▶ **Problem:** **Sampling** often prohibitive (\rightarrow **gsMC***).

* = graduate-student Monte-Carlo method: *Efficient gluing of numerical continuation and a multiple solution method for elliptic PDEs*, CK, Appl. Math. Comput., Vol. 266, pp. 656-674, 2015

Numerical Bifurcation Analysis for Stochastic Systems?

Consider the **stochastic (ordinary) differential equation (SDE)**

$$dx_t = f(x_t; p) dt + \sigma F(x_t; p) dW_t, \quad x_t \in \mathbb{R}^n,$$

$W_t = (W_{1,t}, W_{2,t}, \dots, W_{k,t})^\top$ iid **Brownian motions**,
 $F(x_t; p) \in \mathbb{R}^{n \times k}$; let $\mathcal{D}(x; p) := \sigma^2 F(x; p) F(x; p)^\top$.

- ▶ **Approach 1:** Forward **Monte-Carlo** simulation.
- ▶ **Problem:** **Sampling** often prohibitive (\rightarrow **gsMC**).
- ▶ **Approach 2:** Use probability density $p = p(x, t)$:

$$\frac{\partial p}{\partial t} = - \sum_{i=1}^n \frac{\partial}{\partial x_i} (f(x; p) p) + \frac{\sigma^2}{2} \sum_{i,j=1}^n \frac{\partial}{\partial x_i \partial x_j} (\mathcal{D}_{ij}(x; p) p).$$

- ▶ **Problem:** **High-dimensional** PDE; not even $\sigma = 0$ is easy!

Strategy - Generalization of Continuation to SDEs

Step 1: Recall

$$dx_t = f(x_t; p) dt + \sigma F(x_t; p) dW_t.$$

Step 2: Expand near (locally stable) **deterministic equilibrium** x^*

$$dX_t = A(x^*; p)X_t dt + \sigma F(x^*; p) dW_t$$

where $A(x; p) = (D_x f)(x; p) \in \mathbb{R}^{n \times n}$.

Strategy - Generalization of Continuation to SDEs

Step 1: Recall

$$dx_t = f(x_t; p) dt + \sigma F(x_t; p) dW_t.$$

Step 2: Expand near (locally stable) **deterministic equilibrium** x^*

$$dX_t = A(x^*; p)X_t dt + \sigma F(x^*; p) dW_t$$

where $A(x; p) = (D_x f)(x; p) \in \mathbb{R}^{n \times n}$.

Step 3: The **covariance matrix** $C_t := \text{Cov}(X_t)$ solves

$$\begin{aligned} C_t' &= A(x^*; p)C_t + C_t A(x^*; p)^\top + \sigma^2 F(x^*; p)F(x^*; p)^\top \\ \text{equil. } \Rightarrow 0 &= A(x^*; p)C + CA(x^*; p)^\top + \sigma^2 F(x^*; p)F(x^*; p)^\top \end{aligned}$$

Strategy - Generalization of Continuation to SDEs

Step 1: Recall

$$dx_t = f(x_t; p) dt + \sigma F(x_t; p) dW_t.$$

Step 2: Expand near (locally stable) **deterministic equilibrium** x^*

$$dX_t = A(x^*; p)X_t dt + \sigma F(x^*; p) dW_t$$

where $A(x; p) = (D_x f)(x; p) \in \mathbb{R}^{n \times n}$.

Step 3: The **covariance matrix** $C_t := \text{Cov}(X_t)$ solves

$$\begin{aligned} C_t' &= A(x^*; p)C_t + C_t A(x^*; p)^\top + \sigma^2 F(x^*; p)F(x^*; p)^\top \\ \text{equil. } \Rightarrow 0 &= A(x^*; p)C + CA(x^*; p)^\top + \sigma^2 F(x^*; p)F(x^*; p)^\top \end{aligned}$$

Step 4: Define the **covariance ellipsoid** (“Mahalanobis distance”)

$$B(h) := \left\{ x \in \mathbb{R}^n : (x - x^*)^\top C^{-1}(x - x^*) \leq h^2 \right\}.$$

Covariance Ellipsoids via Continuation

Important observations:

- ▶ Continue the equilibrium $x^* = x^*(p)$ as usual.
- ▶ For covariance ellipsoid one has to solve a **Lyapunov equation**

$$AC + CA^T + BB^T = 0$$

- ▶ During continuation the matrix

$$D_x f(x^*; p) = A(x^*; p) = A$$

is available as a submatrix of $Df(x^*; p)$.

Covariance Ellipsoids via Continuation

Important observations:

- ▶ Continue the equilibrium $x^* = x^*(p)$ as usual.
- ▶ For covariance ellipsoid one has to solve a **Lyapunov equation**

$$AC + CA^\top + BB^\top = 0$$

- ▶ During continuation the matrix

$$D_x f(x^*; p) = A(x^*; p) = A$$

is available as a submatrix of $Df(x^*; p)$.

- ▶ Efficient **iterative methods** for Lyapunov equations exist.
- ▶ A simple **initial guess** for $C(p_{(j+1)})$ at $(x^*(p_{(j+1)}), p_{(j+1)})$ is

$$C(x^*(p_{(j)}); p_{(j)}).$$

Ellipsoids and Distance

Question: What is the distance between ellipsoids?

Ellipsoids and Distance

Question: What is the **distance between ellipsoids**?

Let Q be positive semi-definite then

$$\mathcal{E} := \left\{ x \in \mathbb{R}^n : v^\top x \leq v^\top x^* + (v^\top Q v)^{1/2} \quad \forall v \in \mathbb{R}^n \right\}.$$

defines an **ellipsoid** centered at x^* .

Ellipsoids and Distance

Question: What is the **distance between ellipsoids**?

Let Q be positive semi-definite then

$$\mathcal{E} := \left\{ x \in \mathbb{R}^n : v^\top x \leq v^\top x^* + (v^\top Q v)^{1/2} \quad \forall v \in \mathbb{R}^n \right\}.$$

defines an **ellipsoid** centered at x^* .

Fact: May solve an **optimization problem**

$$\begin{aligned} \delta &= \delta(\mathcal{E}(x_1^*, Q_1), \mathcal{E}(x_2^*, Q_2)) \\ &= \max_{\|v\|=1} \left(v^\top x_1^* - (v^\top Q_1 v)^{1/2} - v^\top x_2^* + (v^\top Q_2 v)^{1/2} \right). \end{aligned}$$

Idea: Use **iterative method** (e.g. SQP) & **initial guess** from continuation to compute δ .

Neural Competition

Consider two neural populations

$$\begin{aligned}x_1' &= -x_1 + S(p - \beta x_2 - g y_1), \\x_2' &= -x_2 + S(p - \beta x_1 - g y_2), \\y_1' &= \varepsilon(x_1 - y_1), \\y_2' &= \varepsilon(x_2 - y_2),\end{aligned}$$

where

- ▶ $x_{1,2}$ = averaged firing rates,
- ▶ $y_{1,2}$ = fatigue/reset variables,
- ▶ $S(u) := \frac{1}{1 + \exp(-r(u - \theta))}$.

Neural Competition

Consider two neural populations

$$\begin{aligned}x_1' &= -x_1 + S(p - \beta x_2 - g y_1), \\x_2' &= -x_2 + S(p - \beta x_1 - g y_2), \\y_1' &= \varepsilon(x_1 - y_1), \\y_2' &= \varepsilon(x_2 - y_2),\end{aligned}$$


where

- ▶ $x_{1,2}$ = averaged firing rates,
- ▶ $y_{1,2}$ = fatigue/reset variables,
- ▶ $S(u) := \frac{1}{1 + \exp(-r(u - \theta))}$.

Look at **noisy fast subsystem** $\varepsilon = 0$

$$\begin{pmatrix} dx_1 \\ dx_2 \end{pmatrix} = \begin{pmatrix} -x_1 + S(p - \beta x_2 - g y_1) \\ -x_2 + S(p - \beta x_1 - g y_2) \end{pmatrix} dt + \sigma^2 F(x) dW_t$$

Numerical Continuation...



For parameter values

$$y_1 = 0.7, \quad y_2 = 0.75, \quad \beta = 1.1, \quad g = 0.5, \quad r = 10, \quad \theta = 0.2.$$

and

$$\sigma^2 F(x^*) F(x^*)^\top = \sigma^2 \begin{pmatrix} 1 & 0.4 \\ 0.4 & 1 \end{pmatrix} \quad \text{for } \sigma^2 = 0.3.$$

Extension to SPDEs

Starting point: (cubic-quintic) Allen-Cahn PDE

$$\frac{\partial u}{\partial t} = \Delta u + 4(pu + u^3 - u^5).$$

$u = u(x, t)$, $x \in \Omega \subset \mathbb{R}^2$, given boundary conditions.

Extension to SPDEs

Starting point: (cubic-quintic) Allen-Cahn PDE

$$\frac{\partial u}{\partial t} = \Delta u + 4(pu + u^3 - u^5).$$

$u = u(x, t)$, $x \in \Omega \subset \mathbb{R}^2$, given boundary conditions.

Main Steps:

1. Compute bifurcation for PDE (e.g. \rightarrow pde2path).

Extension to SPDEs

Starting point: (cubic-quintic) Allen-Cahn PDE

$$\frac{\partial u}{\partial t} = \Delta u + 4(pu + u^3 - u^5) + g(u)\xi.$$

$u = u(x, t)$, $x \in \Omega \subset \mathbb{R}^2$, given boundary conditions.

Main Steps:

1. Compute bifurcation for PDE (e.g. \rightarrow pde2path).
2. Consider the SPDE version (e.g. \rightarrow trace-class noise).
3. Discretize in space (e.g. \rightarrow FDM, FEM, Galerkin).

Extension to SPDEs

Starting point: (cubic-quintic) Allen-Cahn PDE





$$\frac{\partial u}{\partial t} = \Delta u + 4(pu + u^3 - u^5) + g(u)\xi.$$

$u = u(x, t)$, $x \in \Omega \subset \mathbb{R}^2$, given boundary conditions.



Main Steps:

1. Compute bifurcation for PDE (e.g. \rightarrow pde2path).
2. Consider the SPDE version (e.g. \rightarrow trace-class noise).
3. Discretize in space (e.g. \rightarrow FDM, FEM, Galerkin).
4. Apply numerical continuation for SDEs.

PDE: Deterministic Numerical Continuation



SPDE: Stochastic Numerical Continuation



- ▶ **scaling law** of the variance near bifurcation point
- ▶ link to **early-warning signs**
- ▶ Computation on standard desktop computer for SPDEs

Stochastic Continuation - Recent Developments

SODE/SPDE General Framework:

- ▶ “Deterministic continuation of stochastic metastable equilibria via Lyapunov equations and ellipsoids”, **CK**, SIAM Journal on Scientific Computing, Vol. 34, No. 3, pp. A1635-A1658, 2012.
- ▶ “Numerical continuation and SPDE stability for the 2D cubic-quintic Allen-Cahn equation”, **CK**, SIAM/ASA Journal on Uncertainty Quantification, Vol. 3, No. 1, pp. 762-789, 2015.

Climate Science Application:

- ▶ “Continuation of probability density functions using a generalized Lyapunov approach”, S. Baars, J.P. Viebahn, T.E. Mulder, **CK**, F.W. Wubs and H.A. Dijkstra, Journal of Computational Physics, Vol. 336, No. 1, pp. 627643, 2017.

Full Error Analysis:

- ▶ “Combined error estimates for local fluctuations of SPDEs”, **CK** and P. Kürschner, Advances in Computational Mathematics, accepted / to appear, 2020.

Link to Rigorous Proofs:

- ▶ “Rigorous validation of stochastic transition paths”, M. Breden and **CK**, Journal de Mathématiques Pures et Appliquées, Vol. 131, pp. 88-129, 2019.

The Last Slide...

Papers, preprints, etc all available from:

- ▶ **www.multiscale.systems** and **arXiv**

The Last Slide...

Papers, preprints, etc all available from:

- ▶ **www.multiscale.systems** and **arXiv**

Thank you very much for your attention!!